# 5.2 Simplex and Duality

The following linear programs form a primal dual pair:

$$z = \max\{c^T x \mid Ax = b, x \ge 0\}$$
$$w = \min\{b^T y \mid A^T y \ge c\}$$

This means for computing the dual of a standard form LP, we do not have non-negativity constraints for the dual variables.



# Proof

### Primal:

$$\max\{c^{T}x \mid Ax = b, x \ge 0\}$$
  
=  $\max\{c^{T}x \mid Ax \le b, -Ax \le -b, x \ge 0\}$   
=  $\max\{c^{T}x \mid \begin{bmatrix} A \\ -A \end{bmatrix} x \le \begin{bmatrix} b \\ -b \end{bmatrix}, x \ge 0\}$ 

### Dual:

$$\min\{\begin{bmatrix} b^T & -b^T\end{bmatrix} y \mid \begin{bmatrix} A^T & -A^T\end{bmatrix} y \ge c, y \ge 0\}$$
  
= 
$$\min\left\{\begin{bmatrix} b^T & -b^T\end{bmatrix} \cdot \begin{bmatrix} y^+\\ y^- \end{bmatrix} \mid \begin{bmatrix} A^T & -A^T\end{bmatrix} \cdot \begin{bmatrix} y^+\\ y^- \end{bmatrix} \ge c, y^- \ge 0, y^+ \ge 0\right\}$$
  
= 
$$\min\left\{b^T \cdot (y^+ - y^-) \mid A^T \cdot (y^+ - y^-) \ge c, y^- \ge 0, y^+ \ge 0\right\}$$
  
= 
$$\min\left\{b^T y' \mid A^T y' \ge c\right\}$$



# **Proof of Optimality Criterion for Simplex**

Suppose that we have a basic feasible solution with reduced cost

 $\tilde{c} = c^T - c_B^T A_B^{-1} A \le 0$ 

This is equivalent to  $A^T (A_B^{-1})^T c_B \ge c$ 

 $y^{*} = (A_{B}^{-1})^{T} c_{B} \text{ is solution to the dual } \min\{b^{T} y | A^{T} y \ge c\}.$  $b^{T} y^{*} = (A x^{*})^{T} y^{*} = (A_{B} x^{*}_{B})^{T} y^{*}$  $= (A_{B} x^{*}_{B})^{T} (A^{-1}_{B})^{T} c_{B} = (x^{*}_{B})^{T} A^{T}_{B} (A^{-1}_{B})^{T} c_{B}$  $= c^{T} x^{*}$ 

#### Hence, the solution is optimal.

